



JOURNAL

1907

Werner Bucher.







May 27, 1907

Left Washington <sup>at 4 P.M.</sup> on B & O for California via St. Louis & El Paso

The season has been unusually cold and late all over the country.

A few early roses are beginning to blossom & the pionias have just opened at today. Wheat is a few inches high & corn is coming up in a few fields but is not all planted. A few agaves are still seen on cold slopes up toward Harper's Ferry.

Darkness came on before we reached Cumberland.

May 28. A cold night over the mountains. I had to close the window over the screen. A man from Texas who got up early before we were down into Ohio says there was white frost along the way. I slept late and had breakfast as we came into Cincinnati. Here vegetation & crops are just as at Washington & no change was noticed to St. Louis except a little corn big enough to show the rows in southern Ill. St. Louis at dark.



May 29. Reached Kansas City at 7:30 A.M. Vegetation & crops not noticeably different from that of the whole yesterday.

Left Kansas City at 9:55 A.M. on Rock Island for El Paso.

To Topeka the country does not change much but the rich soiled bottom lands give a rich growth to the crops & vegetation not seen before. The rows of corn are slightly more conspicuous than usual but it is rarely 2 inches high. Wheat & oats are thrifty but low. Some winter rye is headed out & a few fields of barley begin to blow heads.

There are lots of good orchards and fine farm homes.

At McPherson the country grows more open & plains like but crops are still good and about the same for condition.

Lots of Jack rabbits (*L. melanotis*) seen after passing McPherson, as many as 7 in one hollow near Pratt.

May 30 - Wake up at Zucumeari,  
and had breakfast before reaching  
Santa Rosa. Raining & ponds full  
of water. Vegetation very late  
grass short but thick, mesquite  
not leaved out, yucca glauca not  
yet in flower but full of buds.  
Verbena, Eriogonum & a few little  
yellow flowers in blossom.  
Gutierrezia getting green.

The same at Santa Rosa &  
over the ridge to Leonida &  
Torreón. Near Torreón they are  
yellow pines on the crests & north slope  
of the mesas and little peaks on both  
sides of the RR and junipers & nut pine  
over the rest of the country except grassy  
flats & valleys. To the west the Joliet  
Mts. are black with timber that seems  
to be yellow pine while still farther  
west the Chaparral Mesa is black  
with timber that seems also to be  
yellow pine. Southeast of Ancho the  
Jicacillos show considerable yellow  
pine along the tops and north slope  
& nut pine & junipers over the sides,  
as does also Carrizozo Peak & range.



About 5-6 miles south of Carrizozo we strike the first trace of Lower Sonoran zone - abundance of Covillea tridentata and Prosopis glandulosa - The yuccas are bearing tillers - more toward radiosa with trunks 1 or 2 feet high. The mesquite is leaving out and a few flowers were seen on the creosote.

At El Zularosa - Alamogorda the vegetation is still backward, and grass has just started. There are almost no flowers, tho at Jirilla the tall Yucca radiosa are full of long spikes of buds and a few are almost in full flower.

Raining and cold nearly all day. Several men wearing overcoats at El Paso.

Left El Paso at about 5:30 and reached Deming just after dark.

Conditions of vegetation the same.

Mesquites in flower at Deming and a few Fouquierias near El Paso in bud but not yet in leaf.

May 31 - Left for Silver City at 10:30  
+ returned at 8 P.M. Got the  
things I would need from my camp  
outfit stored there and made  
arrangements to have my outfit  
kept over for another year.

A heavy frost ~~some~~ 2 weeks ago  
killed the leaves on the walnut trees  
at Silver, but a new crop of leaves  
have come out since. It also  
killed the leaves on the boxelders and  
killed all the fruit for this season,  
even to currants. Since the frost  
vegetation has come on very slowly  
and there is not much green to  
be seen yet and very few flowers.  
Grass is up big enough for good  
grazing and stock is thriving  
several late rains will insure  
abundance of grass this year.

It has been so cold that many  
of the winter birds are still waiting  
in the valley. *Zonotrichias* are  
singing in Silver City & *Cortopos*  
*richardsoni* are loafing around town.  
*Sayornis saya* & *Hirundo* are nesting  
under the porches of the *Depa* hotel at Silver.



June 1. - The 8:30 train was 5 hours late and did not come till 2:30, so we tramped over valley and along dry bed of Rio Mimbres & got a full list of plants & birds & some mammal & reptile notes. The frost of *Yucca radiosa* is sending up great flower stalks that are like asparagus tips only 4 feet long and 2 or 3 inches through. The buds have not yet begun to spread and it will be some time before the flowers are out. The mesquite bushes have a few old blossoms that seem to have been injured by the frost, but are also full of new buds. There are almost no flowers but a few warm days would bring them out in great variety & profusion. The ground is wet below the dry surface.

The Rio Mimbres is as usual a bed of dry sand, but there are many holes where coyotes have been digging down into the wet sand for water. I could not be sure whether they had got any.

The drift along the river has been picked up by 4 men with teams but comes of yellow pines and angles of pine and pine knots are common along the sandy bed. Also little walnuts, pieces of gum and bark from pines along the head of the river.

There are ample evidences of fierce floods at long intervals, but the river bed is too sandy to hold water after the flood has swept by. The underground water is abundant & of excellent quality. Silica City is a forest of windmills and little ranches have started out over the valley in all directions.

The breeding season for birds is scarcely begun here in the valley. One nest with 2 eggs of mourning dove was found, cactus wrens were building, as were barn swallows & song sparrows. The white necked robins are scattered out & probably breeding in the yuccas & we found a family of downy herring owls in a hole in the river bank.



Started for Lordsburg at 2:30  
and colored in zone maps as we  
went. Found nearly the whole  
set of lower Sonoran zone plants  
continuous all the way in the  
valleys except on flats or basin  
bottoms where grass was the  
usual dominant vegetation. On dry  
mesas the *Croton*, *Mesquite*,  
*Ephedra trifurcata?*, *Larrea glutinosa?*,  
and *Yucca schottii* were the conspicuous  
species. At Lordsburg a more  
complete list of lower Sonoran  
species was secured and the  
whole broad valley mapped as lower  
Sonoran. *Croton* bush is the  
dominant shrub over much of dry  
mesa at Lordsburg & up several  
hundred feet above, even on gradual  
north slopes. It is even difficult  
to get many upper Sonoran species  
for the north slopes of the Victoria &  
Pyramid mts., low, bare ranges  
with scarcely any vegetation. I  
merely left the north slopes blank as of  
probably upper Sonoran zone climatic conditions.

Lordsburg - N.M.

Phrynosoma cornuta - One caught  
" woodiata? - ornatissimum?

June 2 - Got a horse & drove south  
about 9 miles to the Pyramid Mts.  
at base of Pyramid Peak, to see if  
there was any evidence of Upper Sonoran  
game species & to look for Amurockia  
& white sided jack rabbits.  
Found no trace of either mammal, the  
Lepus texianus was abundant & I shot  
3 & had time to skin only 2 of them.  
A good lot of big & plant roots  
were secured, but a few scrubby  
junipers near the tops of the peaks  
on north slopes were the only  
real upper sonoran plants found.  
The upper slopes were very barren  
however and had better be colored for  
Upper Sonoran. The warm slopes  
of the mts. show even stranger Lower  
Sonoran elements than the valleys do.  
Fouquieria is abundant & thrifty, the  
big Echinocactus wislizeni is common  
& a tall, long leaved Agave is  
scattered over many of the ridges.  
Several flocks of scaled quail were  
seen with young of various sizes from  
just out to big enough to fly a little.



In P.M. I took the train for Clifton, Arizona and continued N.W. through the same big open Lower Sonoran Valley on a line with Lordsburg to Summit, then down 625 feet lower to the Gila River at Duncan, then down the river to Guthrie (200 feet lower), then over the <sup>3500 ft.</sup> ridge to Clifton on the San Francisco River at only 675 feet above the sea.

The whole trip was in lower Sonoran zone, but along the sides I could in places get approximately the upper limits of the zone.

The change in crossing the Gila valley was less in change of species than in the more advanced conditions of vegetation. Barley was ripe & the first crop of alfalfa had partly been cut & stacked, garden vegetables were being harvested & corn stood over a foot high.

There are good farms & homes along the valley & Duncan is an attractive little town with green fields & big cottonwoods.

Clifton is a big, noisy, booming, banging, slamming, mining town strung along the bottom & side niches of the Frisco canyon for several miles. The mine mills & works are enormous & fill the canyon with a horrid din & vile sulphurous odors & pour down floods of thick gray, poisonous mud into the river. But for the town & mines the canyon would be reeched & wild & picturesque, with steep walls & huge cliffs, numberless caves & caverns & beautiful spots of green & shade in the bottom. The hills are eaten bare by the few burrows of prospectors, but they probably never yielded much grass. I climbed about 500 or 600 feet up just back of town & on the north slope found only *Garryia* as a suggestion of upper Sonoran. The south slopes are pure lower Sonoran to the top, but back 10 or 20 miles to the north I can see what seem to be yellow, vines on the peaks.



June 3. Left Clifton at 7:20 AM. & with  
clear air & no wind got a better distant  
view. Clifton is, by the aneroid, only  
75 feet higher than Guthrie, on the flats,  
but over the ridges between we go  
~~350~~ 360 feet higher than Guthrie.  
The Jila valley is luxuriant in  
plant & animal life and many  
parts are very attractive.

Duncan is the center of agriculture  
& civilization, the abundance of  
water, lands, green fields and  
crops attracting numerous birds.  
There are some pleasant looking  
houses & people & enough of  
the Mexican and wilder element  
to add interest.

Up the dry wash from Duncan  
we got a good view of the Pecos  
Mts. to the west & the glass shows  
quite a lot of Junipers along the  
cold slope. A few Junipers  
were also seen on the slopes down  
near the valley bottom below  
Duncan.

Reached Lordsburg at 10:45 and Hatch  
at 11:45. By the aneroid Hatch

is 250 feet higher than Lordsburg but the country seems level all the way between. It is just a continuation of the plain on which Lordsburg & Deming stand with practically the same set of Lower Sonoran plants all along.

In places the valley is barren or merely grassy & on slopes that tilt to the north there is less evidence of Lower Sonoran, while a south slope will usually be covered with Creosote & Mesquite etc.

The Dry Hatchet Mts. to the south are so high and black with timber toward the tops that I put some in coloring blue above the crest of the range. That too far away recognizes the timber with the glass.

West of Hualapai we go through a low gap in the low Hualapai with plenty of lower sonoran plants going through into the Valley de las Playas, which is a little higher & has partly lower sonoran than the valleys on either side. Then through the gap we



The Peloncillo Range a few quinceas  
come down on cold slopes while  
creosote goes thorough hot  
slopes. Then down about 800  
feet into the big San Simon  
valley which is again pure  
Lower Sonoran across to the  
step base of the Chiricahua  
Mts. Much of the valley is barren  
or grassy but in places  
there is lots of Mesquite,  
Creosote, Yucca schottii,  
Koeberlinia, Ephedra, Larrea  
gutturata? etc.

At the southern end of the valley we  
rise up a long gradual slope  
that soon loses all trace of lower  
Sonoran sparseness mainly grassy  
up to Chiricahua Station  
and over the ridge & down to  
Bernardino Station where abundance  
of Mesquite & lower Sonoran plants  
come in again. The San Bernardino  
valley to the south is a purely Lower Sonoran  
basin that we strike only at corner of.

We next wind through a low pass that  
carries a full set of lower Sonoran plants

around north of College peak and down into  
the great Lower Sonoran Sulphur Springs  
Valley. A few of the big agaves are  
seen in the pass but we are soon  
down in the valley and at Douglas  
at 3970 feet.

Douglas is a big smelting town  
out in the middle of the valley.  
I should guess it had 10000  
inhabitants, well spaced.  
It has a good depot & several  
electric car lines.

The valley is grassy but with  
no visible farming country  
except for the numerous  
windmills & tubs.

Mesquite is the dominant shrub  
of the valley but the whole set  
of Lower Sonoran species occur.  
They also extend over the low gap  
west of the valley to Bisby &  
Haco. A few big agaves &  
some *Dasylianas* occur in the  
gap.

Down in the valley at Haco  
it is pure Lower Sonoran again  
but the sharp little range of



Mts. to the S.W. (The San Joses) have  
a growth of yellow pine along the crest  
& down 1000 feet in the highest cold slopes.

Then we came to the San Pedro  
river valley with its line of  
ranchos, meadows, & cottonwoods  
& the Huachuaca Mts. black with  
timber along the west side.

I can not make out the line between  
the junipers of the lower slopes & the  
yellow pines that grow along the  
crest of the range, but the lower  
Sonoran plants of the valley seem  
to stop short of the wide fans  
that reach out from the canyons.

As we follow north, down the San  
Pedro Valley the river bottoms become  
more heavily covered with mesquite  
and other shrubs and before reaching  
Fairbanks the first *Baccharis bovalis*  
appears & is then abundant &  
conspicuous all along the river flats  
to Benson, where with *Quercus affinis*  
*Atriplex canescens* & mesquite it forms  
a dense cover to much of the valley.

Reached Benson at 6:20 & hunted  
down by river till dark.

Benson, Ariz.

Citellus spilosoma heard  
Am. xerophilus barisi, An old miner  
says they are troublesome around his camp  
in the "Yellowstone Mts." just north of Dragoon  
Summit.

Citellus grammurus - Saw old miner says  
there are gray rock squirrels there too.

Hesperomys hills numerous both  
along river flats & on mesas

Lepus arizonae - common - a few seen.

Lepus texianus - Not common. One  
seen at Fairbaults.

Procyon tracks com. down river.

Lynx " " "

June 4 - In the clear morning light  
I could see no traces of tall pines on  
the Whelston Mts. just west of Benson, tho the  
upper slopes are dark with low timber,  
evidently nut pine & juniper. A few scrubby  
trees appear along the crest of "Yellowstone"  
Mts. just north of Dragoon Summit and an  
old miner working in them says there is  
a little juniper & oak on the north slopes.  
The Dragoon Mts. seem to have only nut pine  
juniper & oaks. On the crest of the  
southern end of the Santa Catalinas  
N.W. of Benson I can see tall timber  
that is evidently yellow pine.

The Sierra El Divisadero show only scattered  
junipers & nut pines over their barren slopes.

Not a giant cactus can be seen  
by sweeping the mesas on all sides  
with the glass, nor have any been  
seen thus far on the trip except  
one in a deer yard here in  
Benson. The nearest I can hear  
of is about 30 miles west of here.

There are none of the Parkinsonias  
or characteristic Atla Valley plants  
here, but the climate seems very  
mild.



Left Benson at 4:30 and soon reached the divide, over which the Lower Sonoran plants pass freely. Then as we descend the first change is seen at Patankito, where the dusky spiny *Cylindropuntia* begins. Then just beyond Patankito a low, stout leaved yucca begins, and immediately on passing Irene the first Giant cactus were seen & soon were numerous on rocky slopes. *Echinocereus wislizenii* also became common and Palaverda (*Parkinsonia microcarpa*) suddenly became abundant. Many of the trees are full of beautiful yellow blossoms. The long leaved Agaves are also common.

To the south of the A.R. the country rises in high ridges between the Whitestone and Santa Rita Mts. and the north slope of this saddle seems to be mainly gussy & probably Upper Sonoran. It is probable however that Lower Sonoran species go through on gulch banks. The Santa Rita Forest Reserve takes in lots of Lower Sonoran zone - What for?

Zueson

June 5. Got a horse and drove out to the Desert Laboratory & met Dr. W. C. Cresson, Prof. Lloyd & Spaulding. Then rode to town & hunted up Herbert Brown & then drove over to Rillito Creek below Ft. Lowell. Then came back to town & got a bicycle and took a bag of traps out near the Rillito Creek & set for *Dipodomys deserti*.

Found giant cactus full of blossoms & green fruit. Saw both yellow & red flowers on different plants of the tree *Opuntia* said to be versicolor - but which I suspect to be two species, versicolor with yellow flowers & solitii with red. *Opuntia spinosa* is also abundant but has the red (purple) flowers. It is loaded with fruit but has a few last flowers. All of the tree cactus are full of nests of cactus wrens, thrashers & workers.



Herbert Brown took me over to the Court house where we found in the basement a few skins of wolves and mountain lions & hundreds of coyotes. The skins are turned in for bounty & then sold by the County treasurer. One tanned skin of a lobo was very brown, but not darker than a beautiful lion *Canis mexicanus* in the park.

a yearling female caught when a puppy 75 miles S.W. of Tucson.

Mr. E. L. Vail, former treasurer at Tucson, who showed us the way & other skins says there are still many wolves on his ranch at the north end of the Santa Rita Mts., east of Tucson. Mr. Brown also reports them seen at Huachuca and killed in the Babagueria Mts.

June 4 - Got a bicycle and went out to traps. got 3 *Dipodomys spectabilis*, 2 baby boys grown, & a *Citellus tereticaudus*. Will catch next train for Casa Grande to try for *Dipodomys deserti* there.

Train was over 3 hours late & we did not leave Tucson until almost dark, so missed seeing the country. Reached Casa Grande late in evening and found a comfortable place to stay at the Woods Hotel, kept by Mr. Shaw.

## Casa Grande Station

June 7 - Took a long tramp over the valley near Casa Grande & saw lots of signs of *Dipodomys deserti*, and *merriami*, and saw 4 *Lepus alleni*, dozens of *L. texensis* & *L. arizonae*. Saw one coyote.

The valley is flat & level, & part of it has been irrigated from the Florence Reservoir, but the water supply gave out and most of the fields have dried up & been abandoned.

The soil of the valley is firm & rich and needs only water to make fine farms. This can be had when the San Carlos Reservoir is put in good condition.

The plants are few in species and colonial in distribution.

For a long distance it will be all creosote, then all mesquites, then all *Atriplex*, then mainly *Yucca* & that is about the limit for the flat country. We found one giant cactus about 6 feet high, but the Mesquites had nearly killed it. Had eaten it full of holes and spiral burrows to near the top.



June 8. Casa Grande Station.

Caught 5 *Dipodomys deserti* &  
2 *sinuatus* & a *Perognathus*.  
got a team and drove S. E. about  
5 miles to the nearest range  
of Mts. to see if there was any  
trace of Upper Sonoran or of  
*Dipodomys spectabilis* but found  
neither. A forest of giant  
cactus encircles the base of the  
mts. & a few reach to the tops,  
common on north slopes, while  
*Fouquieria* & *Polareuda* go up all over  
the mts. *Opuntia spinesa* & *arborescens*  
& *Escobedoia wislizeni* are also  
abundant with the giant cactus,  
as are *Fouquieria*. There seems to  
be no trace of Upper Sonoran in  
these desert ranges, but they are  
all low. I should expect it  
in the Badiguere Mts. to the  
south.

The giant cactus is in flower  
& full of green figs & most holes  
of the Gila Woodpecker.  
A heavy shower of rain & hail  
flooded the roads but missed us.



Cercus giganteus - encircles the foot hills of every range of mts. from where first seen near Irene, east of Tucson, the last point of mountains just before reaching Yuma. See map. On many ranges it struggles up nearly to the tops and it reaches out over stony mesas, but rarely occurs in an open valley or on flat land. It is loaded with flowers & green fruit.

Opuntia spinosa - has almost the same range as the giant cactus around the base of the mts. from Irene to the Mohave Mts.

It is very abundant & very spring is full of fruit and still has a few of its purple or crimson flowers. It is usually full of birds nests.

Opuntia missouriensis - First recognized at Tucson where pale plants had yellow flowers & purple plants have crimson flowers also around foothills at Case Grand Station.

## Case Grand & Yuma

June 9 - Could not get a train to go out to the reservoir so took the belated train at 9 A.M. west.

The country grows dryer & bare - better westward to Yuma.

The mountains are all low & barren & usually black rock, often without visible signs of vegetation. Apparently they are all Lower Sonoran. The Maricopa & Mohave are the highest but just as bare & hot to the tops as any. The only important features of the journey are the beginning & ending of spring.

The soil conditions have a great influence on local distribution of plants and group the species often sharply into societies of moist bottoms, dry bottoms, gravelly valley slopes, stony mesas, stony washes, foothill slopes, rocky gulches, rocky mountain slopes. Certain species of plants happen on each of these formations as it is crossed.



Echinocactus wislizeni - common from Irene to Tucson, Casa Grande, & the Maricopa & Mohawk Mts. Not seen farther west along S.P. Found mainly around the foothills but occasionally one is seen out on flat valley. In blossom at Casa Grande. One cut open & powdered into water which is wet but not good. Tastes like potato juice.

Holocanthos - This thorn tree (Corona christa of the Mexicans) was said to be found about 20 miles south of Casa Grande, but the first place we saw it was about 20 miles west of Casa Grande, in the bottom of the big open valley. It was abundant there and in all the valleys west to the Mohawk Mts. but no farther.

Parkinsonia microcarpa - Palo Verde was first seen just west of Irene, and was there common at Tucson, Casa Grande, and all the way to Yuma <sup>& beyond</sup> in washes and along foothills, but not out on open valley. It grows among

rocks to the tops of desert ranges. Some trees are in flower, others loaded with fruit & others bare of all but green bark.

Iron tree - This lilac flowered tree was first seen west of Mohave & thence all along to Yuma & the Salton Sea. It grows in washes mainly & is often loaded with its lilac colored flowers.

Prosopis glandulosa - From Benson to Tucson & Casa Grande & Maricopa mesquite is abundant and practically continuous. West of Maricopa it is found abundant in places and again not found on dry mesas for long distances. None was seen west of the Mohawk Mts. to Yuma, where only pubescens was seen.

Prosopis pubescens - A few trees of screw bean were seen in the yard at Casa Grande and no more to be recognized until we reached the Gila & Colorado river valleys at Monitor & Yuma, where it is abundant. At Phoenix I think both species <sup>together</sup> are



Acacia constricta - Common from Benson to Tucson. Not found at Casa Grande or west of there.

Mimosa roemeriana? - This catclaw was found at Lordsburg & Duncan New Mex., and at Benson, Tucson, Casa Grande - west to the Mohave Mts., Arizona, but no farther.

Fouquieria spinosa - Common at Lordsburg N.M., at Benson, Ariz., and westward to Tucson, Casa Grande - all the way to Yuma & beyond on mesas & stony slopes.

Chilopsis saligna - Common along the Rillita Creek near Tucson, but not found west of there.

Other plants that range west to Tucson & no farther are:-

*Opuntia engelmannii*,

" " (mushy spinulose)

" " (brown spined)

" *leptocaulis*

Agave fig

*Yucca palmeri?*

" *radiosa*

*Koeberlinia spinosa*

*Ephedra trifurcata?*

There seems to be a marked break in range of species just west of Tucson, but it needs careful working out with camp outfit. It is evidently the change from Lower Sonoran to arid subtropical - or the "lower division" of lower sonoran.

The marginal distribution of giant cactus and palms (Washington) around the sides of valleys would suggest that a hotter belt than is found in the bottoms of the valleys. A series of experiments would show whether this is the case and whether the giant cactus belt is frostless.



Yuma to Indio, Calif.

June 9. At Yuma the Colorado River is out of its banks and has flooded the bottom lands for miles on the west side. Cottonwoods, willows, screwbeans, & thickets of Sarcobatus stand knee deep in water & great lagoons are filled. But after gaining the mesa top the country is dry enough until we reach the Salton Sea near Mammoth. Then we follow the shore of the sea to Mexca on a new road that has been built around it. The old railroad runs through the sea & the line of telegraph poles go out until submerged at the east end & come out again at the west. At Mexca farming begins and the valley is largely cultivated up to Indio. There is a heavy growth of Atriplex, Sarcobatus, & mesquite where the land is not cultivated, but the sides of the valley are still sandy desert of the hottest & driest.

Prosopis pubescens is abundant along the Colorado River at Yuma and reappears in abundance at Mecca & was seen at Coachulla & Indio.

Prosopis glandulosa was not seen west of the Colorado until a few scattered and much dwarfed trees appeared west of Bertram & a few more west of Durand. At Indio it is abundant and large and it reaches along the valley to Coachulla (Coachulla - valley of little shells). Here the pods are full grown & being eaten by Citellus tereticaudus.

Parkinsonia microcarpa - Palatavinda began on the mesa just west of Yuma & was common in washes until we reached the edge of the Salton Sea, then as we followed the shore, it was not seen again until near Mecca, where it is common.

---

"Iron tree", This lilac colored flowered tree was common on the mesa west of the Colorado R. & there all day, (mainly in washes) thence all along to west of Salton, where the last were seen.



## Indio

June 10 Got a horse & rode out to the foothills north of the town, but as I had only a few hours before train time I did not go far enough to get a section down to Upper Sonoran zone. This would have taken a full day, trip back 15 miles to the crest of the mts., for there is only a trace of Upper Sonoran reaching down not over 1000 feet from the summit on the south slope. It is of course more extensive on the north slope. The trees along the crest of the range look with the glass like pinyons & nut pines.

Many of the lower Sonoran desert bushes I could not recall or did not know the names of so I made a rough collection of specimens, wrapped up in paper.

The valley at Indio is fertile, part of it well watered, and in the town well shaded by trees - cottonwoods, palms, mulberry trees etc. Much alfalfa & fruit are raised. Grapes, watermelons & cantaloupes are being shipped.

## To Palm Springs

In P.M. took the train for Palm Springs station, merely a station house on the desert, & then drove over to Dr. William Murray's place at the real Palm Spring - a 6 mile drive across sandy desert with a fierce sand storm raging.

Made part of the trip after dark but saw most of the desert plants seen in crossing the valley north of Indio. Saw also a big wound, a sand dune, covered with a desert willow in bloom.

Could get only a dim view of the palms & other tropical trees at the Springs. Found Mr. & Mrs. Murray a very genial & very interesting old couple. Both are pretty old & feeble & it is hard for them to keep up their place.



## Up San Jacinto Mts.

June 11 - Took a lunch, a canteen of water & an aneroid & started up the steep slope of the Mts. to run a zone line up the first ridge north of ~~Calqued~~ Canyon. Found it steep but easy climbing & pushed on steadily to a little above 4000 feet into the edge of transition zone. Turned back at 4:30 but did not get down the slope before dark & had to make the last 1000 feet very cautiously by feeling. Got back at 9:30 with very weary legs.

The ridge runs mainly east & west and so gives good north & south slopes on opposite sides.

Lower Sonoran zone extends up to about 3000 feet on the north slopes and 4000 feet on south slopes. It is characterized by a scattered growth of *Covillea tridentata*, *Dalia glandulosa*?, *Dalia* "macrocarpa", *Kummerowia canescens*?, *Mimosa coahuilensis*?, *Ephedra antiopphilitica*, *Opuntia echinocarpa*?, *Opuntia crimson flower*, *Echinocactus wislizeni* & *hintoni*.

The slopes are steep & rocky & barren & hot.

Upper Sonoran zone begins on the north slope  
at 3000 feet and extends up to 6000,  
and on the south slope it begins and  
ends 1000 feet higher. It has much more  
vegetation + soil the lower slopes and toward  
its upper limit becomes rather dense  
with chaparral. It is characterized by  
*Juniperus monosperma*, *Pinus monophylla*,  
*Adenostoma sparsifolium*, *Arctostaphylos*  
*vicida*? + *pauciflora*?, *Ceanothus virgatus* + *mollis*,  
*Cercocarpus parvifolius*, *Quercus dumosa* sub,  
*Yucca whipplei*, *Schoenobolus*  
*Prunus andersoni*, *Rhus ovata*, *Philadelphus*  
and others that I collected but  
do not know. It had for birds  
*Aphelocoma*, *Taxostoma*,  
*Pipilo megalonyx*, and for mammals  
*Eutamias merriami* + *Citellus fisheri*.  
Deer tracks were numerous all  
through the zone, but the mountain  
sheep tracks seemed to be mainly below  
in Lower Sonoran. Thoroughbred  
hills were common throughout the  
zone, but I am in doubt as to  
the species. A few Geopline  
ridges were seen + lots of *Neotoma*  
houses among rocks + in bushes.



## Mammals - Palm Springs & San Jacinto Mts.

Ovis nelsoni - Common along the lower slopes of the San Jacintos. Tracks & horns seen.

Odocoileus - Deer tracks are numerous throughout Upper Sonoran zone in going up the Mts. from Palm Springs.

Citellus b. fisheri - Digger squirrels were seen at 3000 feet and heard at 5000 ft on slope of Mts.

Citellus tereticaudus - common in valley and along base of Mts.

Amnospemophilus leucurus - common in valley.

Eutamias merriami, Chipmunks were seen at 3000 and 5000 feet in Upper Sonoran zone.

Neotoma - Woodrat houses were common among rocks & in brushy places all through Upper Sonoran zone.

Thomomys percalanus - Gopher hills are common in the valley at Palm Springs.

Thomomys nigricans? - Gopher hills were occasionally seen throughout Upper Sonoran.

Dipodomys deserti - Common in valley

Dipodomys similis - "

Lepus - Signs of cottontails on Mts.

Sciurus - Runways seen in Upper Sonoran

Transition zone - begins on north slopes at 6000 feet and on south slopes at about 7000, apparently. Tho I did not go above 6200.

Pinus coulteri was the first species reached, but only a few outlying trees before the main edge of tall timber was reached - consisting of Pinus coulteri, ponderosa, and laevis. A few Abies concolor were seen on a steep slope. Quercus chrysolepis and another species of scaly barked oak began about 200 feet lower than the pines & I could not be sure whether they went with Transition or Upper Sonoran.

The line between Upper Sonoran & Transition agrees perfectly at this point with Hall's zone map of the San Jacintos.

## Palm Springs to Los Angeles.

June 12, Very lame & sore & generally used up from 13 hours steady climbing up & down the Mts. yesterday, so I did not get out to do much. At 1 P.M., started back to Palm Spring station against a fierce wind that cut our faces with driving sand & gravel. Took train to Los Angeles & arrived before dark, so saw the country all the way.

Arid Lower Sonoran zone stops rather abruptly at the little station half way between White Water & Banning, and no more *Cercocarpus*, *Atriplex*, *Sarcobatus*, *Franseria*, *Ephedra* or *Echinocactus* were seen. A few *Chilopsis*, *Mimosa roemeriana*?, and a new species of *Cylindropuntia* & the crimson flowered spiny *Opuntia* reach a little further but none of them to Banning.

At Banning & west of there are good crops of fruit, grain, alfalfa, and vegetables, with or without irrigation, and such wild plants as *Eriogonum*, wild oats, grass, weeds etc.



All the rest of the way to Los Angeles the country is largely under cultivation, with many extensive fields, vineyards, & orchards. *Eucalyptus* & pepper trees grow all along & in many places there are palms & farther along great orange groves loaded with ripe fruit.

*Adiantum fasciculata* covers the hills down to edge of valleys and a profusion of vegetation covers all parts of the region.

Meadow larks were common west of Banning & flocks of blackbirds were seen in the meadows & grain fields.

Grain is mostly ripe & much of it cut. Wheat, oats & barley are raised in abundance but largely cut for hay. It was raining just ahead of us & much hay had been wet & some had been spoiled by frequent rains.

Went to the Rosslyn Hotel, a fairly comfortable place.

Sacramento Los Angeles.

June 13 + 14 - Had much writing to catch up and gave maps to carrier. Got mail & wrote letters and laid in supplies for next trip.

June 15, Went out to Pasadena to see Grinnell but found he had gone to the Mts. for the summer. Took train to Oceanside in P.M. & then out to San Marcos where we arrived after dark.

June 16 + 17 - Remained at Twin Oaks over Sunday & Monday, so as to get a supply of cash from the Bank at Escondido. & Got what notes I could but my baggage did not come, so I could do no collecting. Found the valley in flourishing condition with good crops and heavy growth of vegetation covering the whole country. Grains ripe & heavy and were wild oat hay was



can be used. Wheat, oats & barley have given a good crop on mesa land without irrigation. Alfalfa is good on the bottom lands. <sup>Orange, Lemon & grapefruit</sup> trees are loaded with fruit on the mesas & side slopes of the valleys both ripe fruit & the green sets for next years crop. <sup>English Walnuts</sup> Olives show promise of a fair crop. Figs & loquats are bearing profusely. Date & fan palms thrive & many varieties of Eucalyptus are raised for wood. A Camphor tree grows at the Green ranch - Grovillias are abundant & full of flowers.

### Zones -

The bottom of the San Marcos Valley is cold & frosty and evidently belongs to Upper Sonoran zone as do also the north slopes of peaks & the higher south slopes; but the sides of the valley, mesas & lower south slopes & the ridges are unquestionably Lower Sonoran.

The weather has been cool & delightful, with cold nights.

June 18.

Twin Oaks + San Marcos to  
Ocean Side.

Set out traps and hunted all the afternoon - Caught snakes + lizard + Horned toad. Followed a little way up the nearly dry San Luis Rey River + found Alder + a few new cottonwood trees. So many of the bushes are new to me that I shall have to collect them.

June 19 - Caught *Citellus beecheyi*, *Neotoma*  
*paucipes*, *Peromyscus boylii* + *gambeli*,  
*Reithrodontomys longicaudus*, *Microtus californicus*,  
+ *Thomomys b. pallidus*. *Lepus californicus*  
are said to be here + *L. auduboni* is common.  
In evening I went to a rocky gulch  
to shoot bats but saw only one and  
did not get a shot. Killed 2 barn  
owls + saw others.

Found a lot of the strange white  
*Cytisus pulchellus* growing on  
the walls of a canyon near here,  
on the hot slope only, while on  
the cold slope were the more  
abundant but less conspicuous  
*Glycyrrhiza edulis*.



To San Luis Rey, Moosa & Twin Oaks

June 20, Got a team & drove up the San Luis Rey river valley as far as Gopher Canyon, up it to Moosa Canyon, up it to head & over into the Twin Oaks valley & back by Vista & down the Vista river.


Except in the San Marcos Valley found no evident trace of Lower Sonoran unless Eucalyptus, figs & olives indicate Lower Sonoran. No Citrus prints <sup>seen</sup> except in Twin Oaks valley where they do well. The low hills near the coast are mainly covered with wild oats but the hills farther back are covered with dense Chaparral, Adenostoma, Ceanothus, *Persea ilicifolia*, *Heteromeles*, the single leaved *Rhus*, & Oak scrub - and still undivided as to zone limits but are inclined to map all coast exposures as Upper Sonoran & valleys behind mountain ranges as Lower Sonoran. Still many species of Eucalyptus, figs, olives, & english walnuts do well at Oceanside where given shelter of other trees or buildings.

## To Escandido.

June 21. - Engaged a team for several days  
+ drove to Escandido on my way to  
Santa Ysabel + the desert coast by the 4.  
The valley at Escandido seems  
to be well sheltered from coast winds  
and the margins raise good oranges,  
olives, figs etc. The central part of  
the valley is mainly meadows,  
and is evidently cold and frosty  
+ probably belongs to Upper Sonoran.  
The Lower Sonoran belt covers the side  
slopes and runs up the mountains  
without much to mark its upper limit.  
but on south slopes may go to the top  
of the steep, sage covered hills.



To Rawson +

June 22 From Escudido I followed up a valley to the east & south & over a low divide into San Pasquel Valley. This is a wide flat valley with wet bottoms, willow & *Baccharis* flats & meadows and narrow Lower Sonoran borders. The bottom is about 350 feet & raises mainly hay & grain & ~~is~~ <sup>is</sup> ~~upper~~ <sup>sonoran</sup>. All around the edges where most of the ranch houses are, eucalyptus groves are common. Also olive, english walnut, apricot and a few orange groves. The orange trees are full of fruit & look healthy. *Provillea*, *Piper* trees, Umbrella trees, fan & date palms & the big Agaves are common in yards. Several Pomegranate trees were seen full of flowers. At the east end of the valley the Lower Sonoran species are more conspicuous and extensive. Apricot & olive orchards are seen. Also more oranges. Up the Canyon the Upper Sonoran Chaparral begins on north slopes at 700 feet, with a dense growth of *Adenostoma fasciculata*, *Cercocarpus pauciflorus*, *Gutierrezia*, *Ceanothus integrifolius*? C. 

A Menziesia like bush with berries (  
) Rhus (thick leaf) +  
scrub oak (Quercus)

On south slopes Lower Sonoran goes  
to the top of ridges that seem 500 feet higher.  
It is marked by white & blue sage,  
*Eriogonum fasciculatum*, *Dasackia glabra*  
& in places by *Opuntia engelmannii* & by  
the absence of the previously mentioned  
Upper Sonoran species.

~~Here the head of the canyon~~  
A mixture of upper & lower sonoran birds  
characterizes the valley, *Psaltriparus*,  
*Nelson's Oriole*, *Melospiza* etc, with  
*Aphelocoma*, *Melospiza* +

Has the head of the canyon at 1300  
feet large trees of *Quercus agrifolia*?  
& *douglasii*? are abundant. Then  
we cross open grass fields over  
the summit at 1500 feet & down  
to Ramona in an open, mountain  
valley at 1400 feet.

Stopped at the Adams House  
for dinner & was surprised to  
find lemon & orange trees full  
of fruit in the yards. To see olive  
& umbrella & pepper trees flourishing



amid groves of old Eucalyptus, Cypress,  
& pines. There are said to be some  
good orange groves on the sides  
of the valley here, tho the wind is  
too frosty for them.

From Rawona I go east up  
a crooked gulch, then up & up & up  
to a summit at 2500 feet, then  
down a little & through a park  
like valley, then over a ridge &  
down to Witch Creek at 2800 feet.  
After leaving Rawona Valley it is  
Upper Sonoran, chaparral all the way,  
with *Adenostoma fasciculata*, *Arceuthobium*  
*parvifolius*, *Heteromeles arbutifolia*, *Sambucus*  
*glauca*, *Ceanothus divaricatus*, and up  
near 2500 feet a few bushes of a large  
*Arctostaphylos*. White sage & *Artemisia*  
californica run over the top of ridges.

At Witch Creek there is rarely any  
snow or ice, but the summer is  
cool & the zone is Upper Sonoran.  
Still a few fig trees are thrifty & bearing,  
so are english walnuts, pears, peaches,  
apricots, & apples. A small pomegranate  
tree is full of flowers. Grape vineyards  
are common but probably of hardy kinds.

To Santa Ysabel -

At White Creek the chaparral is not so dense or continuous & the big, bald rocky hills are covered with wild oats & scattered live oaks. Not far to the east and southeast are pine covered ridges.

June 23. - A cool night, said to be frost which I doubt, but I slept cold with all the blankets over me I could get.

Continued east over a ridge at 3100 feet & down to Santa Ysabel in a big open valley at 3000. The only farming in this valley seems to be grain & hay. Wild oats cover all the hills & uncultivated land. The valley is evidently straight upper Sonoran, the *Phoradendron flavescens* grows abundantly on the sycamores, willows & live oaks along the creeks.

Along the top of the Volcan Mts. just east of the valley, pines, of which I could make out the long arms of <sup>coastal</sup> ~~Sierran~~ pines, grow all along the coast and of course on the east side. In going over the saddle north of this valley *Quercus californicus* was first seen & is there common. *Otocoris*, *Sturnella* & *Sturniulus* are common in the valley.



## Warner Valley -

Bascanus - Two long red snakes with black heads ran into a gopher hole beside the road. I waited & waited one as he put his head out & found a large horn toad in him.

Phrynosoma blainvilliei - The only horn toad seen was taken from a Bascanus shot on the way down into Warner Valley.

Sceloporus biserratus - The little scaly lizards are common on rocks & trees.

Cnemidophorus stingeri - One was seen & shot at but not secured at the edge of the valley.

## To Warner Valley.

Crossed the divide at 3400 feet and down into Warner Valley to 2700, then across the wide, sandy, open valley to Warner's ranch & up a side valley to the S.E. to Buena Vista, started road ranch & put up for the night.

Found about 100 Indians & Mexicans gathered for a horse race & monte game at the ranch, a rather interesting crowd, men & women betting on the races & joining in the monte game. Just a country Sunday gathering for the only amusement the people have.

Warner Valley is apparently an upper Sonoran valley, the base of any glacial marking vegetation. Grass, fillies, turkey mullein & a low Guthriea like shrub are the valley cover. A few willows & cottonwoods & elder bushes grow along the streams and the chaparral begins well upon the foot hills all around the valley. The chaparral is made up of *Adenostoma fasciculata*, *Adenostoma* <sup>big</sup>, *Ceanothus leucocarpus*, & *canadensis*, *Creoscarpus parvifolius*, *Prunus ilicifolia*, & scrub oak. Live oaks are large & beautiful in groves or singly around the edge of valley.

Warner Valley

Citellus beecheyi. common in & around valley.  
Thomomys abundant all over the valley.

Perodipus, numerous in the sandy valley.

Lepus auduboni - Numerous in the chaparral.  
several seen, one taken.

Taxidea - Badger holes were common  
over the valley, some fresh.

On the Mts. to the N.W. of the valley  
pockets of pine come down half way on  
the cold slopes and pines show along the  
crests of the mts. to the east & Northeast.  
Coniferous timber comes down at least  
1000 feet below the summit of the Volcan  
Mts. just south of the valley

June 24,

Caught a Perodipus, a Microtus  
californicus, 2 Peromyscus bigger & brown  
than boylii + 2 gambeli, shot a pair of  
the redwing blackbirds breeding in the  
tule patches around the springs.

Hitched up + started early, following  
up the eastern arm of the valley to the  
divide towards San Felipe valley.

Was surprised to find Artemisia tridentata  
common in this corner of the valley, also  
Opuntia (like small englemanni) + the herbaceous  
yellow flowered Cylindropuntia (cedarwood?),  
evident overappings from the desert.

Crossed the divide at 3450  
+ descended rapidly into the San  
Felipe valley. At about 2700 feet began  
to strike traces of Lower Sonoran  
in Prosopis glandulosa + Mimosa  
roemeriana, Chilopsis + Opuntia b. canadensis



## San Felipe Valley

Citellus beecheyi common  
Citellus tucumanensis, One seen.

Amaspermophilus lineatus, common  
many seen but none secured.

Neotoma \_\_\_\_\_, housed common  
both in rocks & in bushes  
& cactus.

Lepus arizonae? Cottontails are  
very numerous. I shot 3 & saw  
at least 3 dozen. They were  
warty among the rocks.

Lepus texensis deserticola, abundant  
Four shot & dozens seen.  
Found among mesquite & creosote  
bushes.

Thomomys hills are numerous.

Microtus californicus - Signs seen along creek.

Dipodomys agilis? Holes & trails  
of a kangaroo rat of this size  
& habits of Agilis were numerous.

Canis - A coyote skull  
was picked up.

Scapanus - A mole ridge was  
seen at Banner in the Canyon.

## San Felipe Valley

The whole San Felipe valley below the  
2400 foot contour and up to at least 3000  
on foot slopes east of valley should be  
mapped for Lower Sonoran. Abundance  
of Mesquite (P. glandulosa) covers the  
whole valley and at the eastern end  
below the 2400 foot contour Creosote  
bush is abundant. Other desert  
Lower Sonoran shrubs are Neonotia  
roemeriana, Lycium arborescens? & "sp. shrub",  
Ephedra, Thamnosma, Coleogyne,  
Atriplex polycarpa?, & canescens, Cordia?  
, Opuntia engelmannii & schinocarpa?  
Eriogonum engelmannii, 3 or 4 composite  
shrubs & Baccharis borealis.

The lower end of the valley  
opens out & down onto the desert  
& Salton Sea.

Upper Sonoran zone comes  
down to the edge of the valley on the  
west & south. On the west it is  
mountain chaparral, on the south  
& east desert species such as  
Juniperus monosperma, Yucca  
wahavisensis? Agave (medium),  
mixed with the mountain species.  
Junipers cover the north slopes & easts.

of the barren desert ranges east  
+ south of the valley.

The main mountain slope west of  
the valley is densely covered with  
chopwood nearly to the top, or up  
to the pines + firs that cover the  
top + north slopes down the canyons  
1000 feet or so. This chopwood  
is especially dense all along the  
road from Banner to the  
summit of the range near Julien  
or from 2600 to 4100 feet.

It is composed of *Adiantum fasciculatum*,  
*Canathus diversicatus*, - (spiny leaves)  
and (wide soft leaf), of *Prunus*  
*ilicifolia*, *Heteromella*, *Rhus* (big leaf)  
*Arctostaphylos patula*?, *Cercocarpus parvi-*  
*folius*, *Ribes cereum* (common), *Cercis*  
, *Arctostaphylos* (yellow),  
*Lonicera* (white), much *Quercus*  
*damosa*, <sup>hypericoides and</sup> many less conspicuous  
plants. In the gulches down at  
Banner, *Populus fremonti*, *Platanus*,  
*Alnus*, *Salix nigra*, *Sambucus glauca*,  
*Vitis*, *Aureus agrifolia* + *englensis* are  
abundant.



Eutamias merriami, heard a chipmunk  
first at the top of the Banner grade  
& a couple of miles east of Julian.  
A miner at Banner told me there were  
chipmunks up in the pines.

Atell's beecheyi - One seen just below  
Julian & others a little lower.

Microtus californicus - Microtus umoyi  
were seen in the meadow just  
below the edge of town.

Thomomys hills are common  
all over the mts,

Near the top of the ridge, Quercus  
californicus - Chrysolopsis biguttatus <sup>at 4000 feet</sup> just  
below the Pinus coulteri are most.

The chaparral <sup>scattered</sup> reaches nearly summit  
& open pines & oak woods covers the  
top of the ridge. The big <sup>green grassy</sup> ~~open~~ ~~valley~~  
~~valley~~ <sup>valley</sup> Arctostaphylos begins on the  
west slope in scattered bunches.

Reached Julian in time for  
supper, then I stopped & hauled at  
Banner & fed my horses & skins  
a cottonlaine. Could see no other  
pines than coulteri.

June 25.

Had to stop & get 3 shoes put on  
my horses, then after going a mile  
or two one of the bars of the pole broke  
& I had to tie it up & return & have  
it welded, so did not finally get  
off until after 10. Came SW. from  
Julian, keeping near 4000 feet for  
a or 8 miles through half open,  
black oak & coulteri pine forests.  
Tongues of upper Sonoran chaparral  
come up on both slopes but  
most of the country above 4000 is  
transition zone. It is open

& grassy & characterized as much  
by absence of chaparral as by its  
zone marking species. Besides  
*Pinus coulteri* & *Quercus californica*  
there are scattered bunches of a big  
smooth leaved *Arctostaphylos*,  
*Rhamnus californica*, *Symphoricarpos*  
*araphius*, *Wyethia* with woolly  
leaves the size of my hand, a woolly  
leafed *Thermopsis*, a big pink purple  
*Lupinus*, wild roses and in  
shady places patches of *Pteris*.

The little farms are mainly meadows  
with orchards of thrifty apple trees  
full of young fruit & some potato  
& corn patches. The Wyola  
valley, just west of Julien is  
famous for its cherries, both  
in abundance & quality.

The road began to go down  
a long grade into the San Diego  
River Canyon & for about 2000  
feet was in slopes covered  
chaparral of the standard species,  
largely *Adenostoma* & *Ceanothus*  
& scrub oak, but with  
*Rhus*, *Heteromeles*, *Prunus ilicifolia*



*Rhus trilobata* + the single leaved <sup>*Rhus laurina*</sup> kind  
+ many others. Scattered live oak,  
*Quercus agrifolia* + *densa*, are the  
principal timber.

Just below the falls, <sup>at about 2000 ft</sup> the chaparral  
ceases on hot slopes & is replaced  
by *Hosackia glabra*, white sage &  
herbaceous plants. At the bottom of  
the canyon at 1000 feet *Yucca*  
*Sonorae* Chaparral comes down on  
cold slopes, but the hot slopes are  
more open & barren with a few lower  
*Sonorae* species of plants.

*Mimosa roemeria* is abundant  
along this part of the canyon on  
the hottest slopes, but there  
are no crops & not many  
lower *Sonorae* plants.

Along the river an abundance of  
*Populus fremonti*, *Sycamore*,  
*Alnus*, *Salix nigra* & a  
yellow willow & in places  
*Baccharis borealis* & a fine leaved  
species. Elderberry bushes  
are full of berries that are here getting  
ripe & hundreds of *Phainopepla*  
are feeding on them.

## To Lakeside.

The Indians raise a little wheat & barley & squashes & corn but nothing to mark a zone. The benches & mesas would doubtless raise good oranges & other Lower Sonoran products. The soil is good & there is plenty of water at present in the San Diego River besides that in the large flume bordering the valley. The great abundance of elderberries, now ripening, may have attracted the Indians as well as the Spaniards to this valley. After turning west the valley is narrow and canyon like and the sides are covered with chapparal. The moist ocean winds come through it & evidently cool it to the an upper Sonoran temperature, while the valley above is cut off from this wind. About 5 miles before I reach Lakeside the valley opens out with wide bottoms & some ranches. Around the slightly elevated margins of the valley the eucalyptus, pepper trees, and orange groves would indicate Lower Sonoran zone. Even at Lakeside, near the lake & flat



## Lakeside

The eucalyptus, Washingtonia palm, Pepper trees, umbrella trees, Ohauas & big Agave grow to perfection while along the slopes to the south & west are extensive & beautiful groves of Oranges & Olives, of old, healthy, bearing trees. The whole valley except the damp bottom should be mapped for Lower Sonoran. It is sheltered from the ocean winds by the mountains to the west, is rather dry & hot.

June 24. Got an early start from Lakeside, and driving due west about 5 miles, turned north & followed up Eyecanon Cr. a few miles, then over the ridges by a new & easy grade to the Summit & down into Paway Valley, then over a low ridge & down to the Bernardino River & then slightly up & over to Escondido.

The sides of the valley west of Lakeside & of Eyecanon Creek valley are Lower Sonoran, with little to mark them except absence of most of the species of Upper Sonoran Chaparral, and its place filled by *Opuntia engelmannii*, white sage,

Labioid + Poway Valley.

*Hosackia globosa*, wild oats and small plants, turkey mullen, a slender *Ocotilla*, and others. The introduced species such as Oranges, Eucalyptus, pepper trees, olives, big Agaves, Grapes, Washingtonia palms, Umbrella trees, & the Oleander certainly indicate Lower Sonoran zone.

A peculiar feature in zone distribution here is that Lower Sonoran runs highest on S.E. slopes, the slope sheltered from cool ocean winds and exposed to the hot sun.

In the bottom of the San Diego River valley there are 3 species of *Baccharis* or *Beckaria* like shrubs, many willows & especially the narrow leaved *S. rigens* or form of it, some sycamores and numerous cottonwoods (*P. fremontii*). It is doubtful if the bottom of the valley can be considered Lower Sonoran or at least the waist part of it.

Poway Valley lacks most of the Upper Sonoran species of clapped over its warm slopes but has Eucalyptus groves, Gravel trees, Pepper trees & the big Agave marginals.



Bervardo Valley  
to  
Eseondido

Then over a low ridge to Bervardo Valley. The warm slopes & sides of the valleys are all Lower Sonoran open & hot and dry, well sheltered by mountains to the west. Eucalyptus & pepper trees were all I could see of Lower Sonoran growth except one much neglected but fairly good Orange grove.

From here over to Eseondido the Lower Sonoran open slopes seem to be continuous on the south east slopes at least.

Coming into the valley at Eseondido from a new direction seems to add more evidence of its Lower Sonoran affinities. Orange & grape fruit & lemon groves are more numerous & extensive than I thought & many other trees and plants, palms, agaves, & such are abundant.


Reached Twin Oaks at 6 P.M. & staid over night.

Oceanside

June 27 Drove to Oceanside, & got mail including *Microtus* bulletin. mss. to be gone over, a two days job. With this & reports & express account, packing specimens & writing letters I could not get off until July 2.

July 2 - Oceanside to San Diego.





Santa Ana to San Francisco

July 4 - Got up at 5:30 at Santa Ana & reached Los Angeles at 7. Left for San Francisco by Coastline S.P. for San Francisco.

From Santa Ana to Los Angeles is level, fertile, highly cultivated valley country or extensive hay & grain fields. Great orange groves of beautiful trees loaded with fruit (part of them) are seen in many places. English walnut groves are numerous, healthy & full of fruit. Vineyards, peach & apricot orchards are common. Eucalyptus groves & rows of palms & pepper trees are seen all along. There is no chaparral. The whole valley is intense Lower Sonoran with no marked trace of Upper.

Leaving Los Angeles we follow up the river valley to Fresno & Burbank & ~~Fresno~~ <sup>Chico</sup>, a wide, flat valley, evidently rather dry, often sandy & desert like with much cactus (*O. engelmannii*? & *chinacarpa*?), with white sage & *Hosackia glabra* & several forms of *Baccharis* & other composite shrubs.



\* more weeds (mustard, radish,  
Toadstole thistle, etc) than native plants.  
An occasional bush of Chinese grows on  
hawthorn & elder along bottoms.

Vineyards are numerous & extensive,  
there are large olive & Walnut orchards,  
some orange groves. Lots of Eucalyptus,  
rows of palms & pepper trees and  
great fields of grain (wheat, oats & barley)  
towards ~~Chatworth~~ <sup>Chatworth</sup> the soil is better &  
more grain & pasture & hay are raised in  
the valley, but more olives, apricots,  
& orchards along the foothills.

The mountains north & south  
of Burbank are covered with chaparral,  
in places dense chinquapin & scattered  
oaks & Rhus & hetttranchs.

While the valley is extreme lower  
Serran the mts. are intricately dissected  
with lower & upper according to  
exposure & elevation.

After passing Chatworth we go through  
several long tunnels under a rocky  
ridge & come out into an open valley  
at Santa Susana, where abundance of  
*Quercus lobata* was first seen.  
The valley is mostly open & bare of

of chapparal. Grain, olives, Eucalyptus, apricots, alfalfa & beans are the main products. Even the hills are covered with sage, *Hesokia*, *Artemisia californica*, & a few *Quercus agrifolia* & could be cropped as Lower Sonoran.

Only the distant ranges north & south show dense chapparal of upper Sonoran.

Continuing westward down open valleys the country is similar, but more fertile & better cultivated on to Oxnard.

Besides grain & occasional orchards there are extensive fields of beans & between San Luis & Oxnard thousands of acres of sugarbeets. Also much alfalfa & numerous groves of large eucalyptus. The mountains have disappeared on the south but are high far to the north.

A cool breeze comes in from the coast & vegetation takes on a ~~more~~ fresher & greener appearance.

Oxnard is a ~~wide~~ town of small houses full of trees, eucalyptus, pines.

We soon ~~cross~~ the river & go through more & bigger fields of sugar beets & beans, before reaching the coast at Ventura.



West of Ventura the road is cut into seen  
flaps that is steep & cool & chappal  
covered & seems to be upper Sonoran.

Then asphalt & oil works are passed &  
a little before reaching Santa Barbara the  
coast flattens out for a space & the  
chappal is less & ~~now~~ some crops &  
trees are raised. At Santa Barbara there  
are abundance of Eucalyptus, pepper trees,  
figs, olives, a few orange trees in yards,  
rows of fan & date like palms, the  
yuccas, *Morisona glauca*, clauthus,  
a few rubber & magnolia trees, orange, loquats,

Tall trees appear above the black  
chappal slopes of the Santa Ynez Mts.  
to the north - just a few along the crests.

West to Point Conception this  
range strip of semiarid Lower Sonoran  
continues, each little point & over  
ridge serving to cut off the cool  
west wind. Little towns & settlements  
are scattered along but grapes  
are the principal crops & there is  
little to mark Lower Sonoran except  
absence of Upper Sonoran chappal.

*Artimisia californica*  
*Ambrosia*

*Baccharis "dumifolia"*

*Lupinus* (shrubby blue)

*Sedum sedula*?

*Eriogonum fasciculatum* "prostratum"

*Samolus*?

Blue sage

abu.

"

"

"

"

"

"

"

At Point Conception, or more markedly at Arquello point, the surface changes to high, rocky shore & steep bluffs, & the vegetation to a dense cover of low wind beaten shrubs & plants as noted on the opposite page.

Big surf comes rolling in the coast is picturesque & attractive. The air is much cooler. Further inland to the east, up the Santa Ynez & Jesus Maria valleys I can see yellow slopes that suggest big lower Sonoran valleys, warmed up by distance & sheltering hills. But the coast strip continues, what I assume to be Upper Sonoran to where we leave the coast & follow up the river valley & beyond as far as I can see up the coast.

As soon as we get back behind the first row of hills from the coast all shrubs cease & there is only wild oats & a few scattered & rather scrubby *Quercus agrifolia* - grain & beans are the only crops until we come down into the big valley of Sta. Maria. This valley is largely open to the coast



and while the western edge of it along the railroad is tinged with chaparral & raised mainly sugar beets, potatoes & barley the eastern and greater part are yellow with grain & stubble & wild oats & dotted with live oaks, eucalyptus groves & orchards of some kind. A few scrubby little palms & pepper trees suggest that farther back the valley should be mapped as Lower Sonoran. Yellow oak covered hills east of the valley also support this view.

Along Arroyo Grande, near Ocean, are extensive marshes, of weeds & willows & some tulips & cat tails, full of birds, mainly blackbirds. A few small fan palms & fine groves of eucalyptus grow at Ocean, but the coast chaparral, as noted on last page with Rhus, Sambucus & willows added predominates to Groves as we follow the coast. At Groves we go up a narrow willow & Sycamore gulch, with some live oaks & wild oat slopes & some chaparral slopes, a mixture of zones.

## San Luis Obispo

At Edna another beautiful little island valley opens up, well sheltered from the sea winds by a steep little range of mts. The native plants are mainly live oaks & wild oats & dense slopes of chaparral on some of the hills to the west & here there a patch of wheat. Grain & beans are the principal crops but eucalyptus groves are common as well as pines & cypresses around the houses. At the foot end of this valley San Luis Obispo lies in between the hills. In the town are small palms (palm & date like), pepper trees, olive, fig & peach, english walnut, apricot, a few good thistles, oranges, grapes, & lots of eucalyptus, pines, sycamores & live oaks.

Barley, oats, & wheat are the principal crops raised from fruit.

From San Luis Obispo we climb rapidly to the crest of the coast range over high slopes of wild oats on south & east & chaparral on unprotected west slopes. There are live oaks & sycamores with the wild oats and the chaparral is *Artemisia californica*, *Gutierrezia*, blue sage, *Adenostoma*.



Over Mts. to Santa Margarita

*Eriogonum fasciculatum*, *Yucca whipplei*,  
*Amelanchier* & many other shrubs  
& could not be sure of. ✓

Fire is running over the mountain  
sides & miles of wild oats & some  
chopped are burned & burning.  
After going through many tunnels we  
go through a long one & come out  
in a high mountain valley with  
dense chopped around the sides  
& lots of *Pinus sabiniana* (if I know it).  
The sheltered slopes of this valley are  
open, wild cat farms, full of big  
*Quercus lobata* & *agubia*.

We now go down to the north about  
4 miles from tunnel to Santa Margarita  
(935 feet). Cottonwood, willows &  
sycamores along stream. *Q. lobata* &  
*agubia* along bottom, *Pinus sabiniana*  
= chopped along sides of valley.  
Cat bay is the only crops. With some  
big fields of potatoes, alfalfa, & garden  
vegetables, small orchards. Poor  
farming. At Temblon a good  
olive orchard & broad valley  
full of *Q. lobata* suggest some Spanish

A few neglected orange trees in a wedy  
yard cling to existence.

No more pines in sight. At San  
Miguel a few pepper trees are seen  
but there is very little farming & that  
is of the worst kind. All arroyos  
are full of grass & weeds & most are  
drying. Cattle & hay are the  
chief industry. Still the lower  
Sorean part of the valley widens  
& great parks or calling meadows  
are all wild oats & big oaks.

To Kings City & an hour beyond  
when it got dark, the valley is  
similar, thinly settled, mainly  
hay & grazing. The little  
farming is poor & carelessly  
done. Upper Sorean is capped  
covers the ranges & the west  
with tall pines often showing  
along the crests of the higher  
ridges. We run so close  
under the foothills on the east  
as to hide all beyond except  
now & then a glimpse.



## San Francisco to Capistrano

July 5. Got into San Francisco about 2 A.M. but went to bed & slept till morning on the train.

Found Dr. Merriam and used most of 3 days in laying plans for the summer's work.

July 7. Started south at 5 P.M., tried to go by the Valley route to Los Angeles but every berth was taken on the night train so I had to go back by the Coast line.

Went from Oakland to San Jose where I caught the down train & in this way saw the strip of country east of the bay.

July 8 - got into Los Angeles after noon & too late to get a train out so remained & hunted up botanics & maps

July 9. Went down to Capistrano got out my baggage & set a line of traps

Capistrano -

July 9 & 14 - Trapping & collecting at  
Capistrano -

July 15 - Drove up Trabuco Canyon  
into the Santa Ana Mts. to 2300  
feet. Entered the mountain canyon at  
1300 feet where upper Sonoran species  
cover north slopes & followed up beautiful  
wooded creek bottom about 5 or 6 miles  
to near end of wagon road - to the first  
grove of Pseudotsuga macrocarpa at  
2300 feet. There were other scattered  
trees & groves of it higher up, but  
no amount of timber anywhere in  
sight on the mts. We could see 3 small  
groves of pine along the crest of the range  
& Forest Ranger Robbins said it was  
coulter pine, as it looked to be. He said  
there were no other pines on the mts.  
but reported a white oak (probably englemanni)  
higher up and the golden cup oak (chrysolepis)  
and said there were a few californica laurel  
(umbellularia) in the canyon. He thinks there  
are some junipers over by Hot Spring, but  
is not quite sure. For lists of chaparral  
see note book.



July 18. To Santa Ana.

July 20 - Drove up Santiago Canyon to mouth of Silverado Canyon about 1000 feet and some 15 miles east of Santa Ana. Drove through miles of beautiful bearing Orange & Lemon groves with enough eucalyptus trees along the roads to furnish fire wood for the ranches. The whole valley to Orange & beyond & to the foot hills is laid off in squares of streets and fruit trees.

There are numerous fields of english walnuts, some grapes and out on the foothills olive orchards.

Palms, of many kinds, and pippertrees are numerous and pomegranates, Loquats, pears & apricots are raised.

Over the valley there is scarcely a native plant left, but *Eriogonum setigerum*, *Croton californicus*, *Medea elaeagnifolia*, *Grevillea* & other weeds fill the waste places. Along the foothills cactus is abundant, mainly the 3 kinds of prickly pear and a few *O. bismarckiana*. *Hosackia glabra*, *Antennaria californica* & *Eriogonum fasciculatum* & white sage (*Quercus laevis*) grow on the foothills.

In the low, open part of Santiago Canyon there are abundance of sycamores, willows, *Baccharis viminea*, and live oaks (*agrippa*). At 1000 feet on the canyon slopes, or at least the cold slopes, the real chaparral begins in *Rhus laurina* + *ovata* + *tillobota*, *Rhamnus crocea* + *californica* + *toментilla*, *Cercocarpus pauciflorus*, *Quercus dumosa*, *Aleostoma fasciculata* etc.

From here the mts. are densely covered with chaparral to the tops, with only 2 small patches of Douglas spruce (*massachusetts*) on cold slopes near the summits as shown by field glass.

There is no farming in the canyon. The lower part is held as a park for Santa Ana + is full of campers. Above are some mines in Silverado Canyon + loads of ore were met.

The canyon is open for a long distance + beautiful + clean with big sycamores + live oaks. A good stream comes down through it.

On the return trip we drove around through Orange to Santa Ana.



A Woman living on a ranch  
7 miles south of Santa Ana and  
somewhat higher up, on the foothills  
says she has kept tomatoes and  
chile plants over winter for four  
years before they were killed by frost.  
As they are killed by the slightest  
frost this is a significant fact.  
She also says sweet potatoes  
& peanuts are extensively raised on  
the dry, unirrigated uplands,  
also beans, lima, navy & black-eyed.  
We saw miles of beanfields

June 21. Santa Ana to Newport.

Sunday afternoon, after finishing reports  
& maps & plant lists we took train  
down to Newport on the coast.  
After passing the fruit ranches we  
came through great fields of beans  
and grain, wheat, oats, barley & corn,  
then over wild oat and grass lands  
to the shore. There are some washes  
& ponds by the way & a lot of  
little washes & some slits at the  
ponds. The soil is sandy toward  
the shore & all tracked up with  
*Perodipus* & other little tracks.

There is little vegetation of  
native species, but turkey vulture  
*noctua* & *grioides* suggest Lower  
Sonoran. *Opuntia occidentalis*  
grows on the sides of gulches near  
the coast. The usual beach plants are

The San Joaquin hills to the  
south are covered with mainly  
wild oats & mustard, but patches  
of low chaparral appear to be of  
*Artemisia californica* & *Eriogonum*  
*fasciculatum*, neither of which  
mark a zone.

## To Corona.

July 22. Left Santa Ana on 11:45 train for Corona. Passed through the same kind of orange, lemon & walnut farms to Olive. Then after crossing the Santa Ana River came into a desert strip of cactus, sand & waste land. It is old river flats, stony & sandy - poor soil but rich in animal life. The sand is all tracked up with small things - *Perodipus* etc - & would probably yield several desert species. It could be worked from the little town of Olive by walking a mile north.

We then follow up the Santa Ana River through the canyon to Corona, with lower Sonoran plants all along the valley including cactus, *Eurotia setigera*, *Baccharis viminea* etc.

Upper Sonoran Chaparral comes down near the valley edge on slopes south of the river and covers the Santa Ana mts. densely to the tops. Half a dozen little patches of big cone *Pseudotsuga* appear in the upper ends of N.E. canyons above 2000 feet. Most of the trees seem small.



The hills north of the Santa Ana R. are generally low & bare of even chaparral. Patches of *Artemisia californica* & *Eriogonum fasciculatum* cover some of the upper slopes but have little zonal significance. From Corona the higher cold slopes have a suggestion of some heavier chaparral, but most of these hills may be considered Lower Sonoran.

The country about Corona is hotter and dryer Lower Sonoran than about Santa Ana. Oranges, lemons, grapefruit, olives & pomegranates are the principal crops back to the foothills - to about 1200 feet.

In the afternoon we drove up the canyon south of Corona, up into the dense Upper Sonoran chaparral of *Rhus laurina* & *obata*, *Ceanothus crassifolius*, *Greocarpus*, *Quercus dumosa*, *Rhamnus californica*, *Tamulus*, & *crocea*, *Adenostoma fasciculatum* etc. Could see several small groves of Douglas spruce (*Pinus jeffreyi*) on NE slopes far above, but the heavy chaparral runs to the top of the range.

## To Coldwater Canyon.

July 23 Drove from Corona about 10 miles south to Temescal and up into the mouth of Coldwater Canyon and stopped at the Glen Ivy Hotel. This is at 1400 feet at the mouth of the canyon and lower edge of chaparral. The note says stop & bushy base of the house. The canyon cuts in narrow & steep sided & a tangle of trees, bushes & vines. A fine cold, clear creek comes out of the canyon & is at once harnessed in pipes & flumes & set to work in orange <sup>& lemon</sup> groves - from here to Corona and then furnishes part of the water supply for the town of Corona.

A hot sulphur spring also comes out just below the mouth of the canyon & is used for a big bath house below the hotel.

The valley is semi-arid with lots of cactus & some scattered chaparral & iron oaks. There are grain fields and down along Temescal Cr. some alfalfa fields and on the benches a few orange & lemon groves. At Glen Ivy Hotel there is a good bearing orange grove of several acres & lots of fine large fan palms.



The Canyon is narrow & bushy  
& full of live oaks, sycamores, alders,  
willows & farther up maple & some  
*Pseudotsuga macrocarpa*. The chaparral covers  
the north slopes abundantly & the south slopes  
sparingly to the mountain tops.  
The conspicuous species are *Rhus laurina*,  
*crata*, *Quercus*, *trilobata*, *Rhamnus*  
*californica*, *tomentosa*, *ilicifolia* (and  
*crocea* in the valley), *Ceanothus*,  
*Arceuthobium parvifolium*, *Ostrya*, *Arctostaphylos*,  
*Rhamnus*,  
*polytrichum* & *dactyloides*, *Artemisia*,  
*californica*, *Eriogonum fasciculatum*,  
*Yucca whipplei* etc. It is so  
dense that travelling off the trails  
is almost impossible, and the  
the only trail does not go far  
up the canyon. Even on it the  
grapevines & poison ivy make a  
tangle difficult to penetrate.

Mammals are scarce in the  
canyon except *Neotomas* & 3  
species of *Peromyscus*, *Sciurus*  
& gray foxes. There are no signs  
of *Thomomys* in the canyon & only  
*Microtus californicus* at the lower end.

Just below the hotel the valley is dry  
& sandy and *Perodipus* and the cactus  
wood rat & *Thomomys pallidus* are  
abundant, also *Lepus auduboni*.  
All of the Lower Sonoran plants of the interior  
valley country are common - Cactus, *Baccharis*  
*virinea* & the leafless species, *Eremocarpus*  
*retigonus* & *Croton Californica*, *Cucurbita foetidisima*,  
*Madia elegans*, *Senecio doglasi*, & *Biscutella chrysantha*.  
Lower Sonoran runs up against the mountains  
& on steep south slopes keeps out the dense  
chaparral. Oranges & lemons & palms do  
well, even when the lower edge of chaparral  
is cleared out to make room for them.

The hills east of the valley look  
bare and brown, but are really covered  
with a tangle of white & blue sage,  
*Eriogonum fasciculatum* & *Artemisia californica*.  
There are scattered bunches of *Rhus glabra*  
and a few other bushes in the gulches and  
a trace of heavier chaparral can be seen  
at the edge of cold slopes. All but the  
higher cold slopes of this low, rough  
plateau, or group of hills should be mapped  
as Lower Sonoran. It is apparently  
used for stock range & bee pasture.



Zemereal to Elsinore.

July 24.

Left Glen Roy Hotel at 2 P.M. and drove to Elsinore, 14 miles south east and put up at the Elsinore Hotel.

Followed up a narrow and often rough valley from Zemereal to Elsinore valley with the Santa Ana Mts. rising black & chaparral covered on the west and brown "sage" & "buckwheat" (*Ramaria* & *Artemisia* & *Eriogonum fasciculatum*) covered hills on the east. More Douglas spruces (*macrocarpa*) and Coulter pine appear in the upper canyons than I have seen before and as usual the spruce comes down to about 2500 feet on cold slopes while the pine appears along the higher part of the range.

The bottom of the valley & the south slopes of hills to the east are Lower Sonoran with the same set of plants seen in the valley at Zemereal. Many bee ranches are seen in corners of the valley and some orange, lemon & olive groves along the foothills to the west.

July 27+8

Elsivore.

at Elsivore the valley is open, with low hills on the east & the Santa Ana Mts. rising ~~abruptly~~ from the west side of the lake. The lake is 3 miles wide & 7 long, and is said to have risen 20 feet in the last 2 years. but is still 8 feet below the overflow level.

Much of the shore is flat & grassy and is used for pasture or hay. There are few plants to indicate the zone, so I have mapped it as upper Sonoran.

Just back from the shore are abundant *Pucca bonalis*, & *Baccharis viminea*, also a few *Baccharis emoryi*, some *Opuntia occidentalis* & *bernardina* - *Eriogonum* *sitigenum* & *Croton californicus*. On the evidence of their presence and absence of upper Sonoran species I have mapped the valley & foothills as Lower Sonoran, including a foothill strip along the west side of the lake below the chaparral that is largely cultivated and occupied by orange, lemon & olive groves. In town pepper trees, Eucalyptus, Palms, Oranges, olives, acacias, and big agaves are common, but the crops out over the valley are mainly grain or hay.



For the three mornings at Elsinore the fog covered the Santa Ana Mts. to the base but soon lifted after 8 or 9 o'clock. It sometimes even hides the lake and the hills to the east but the valley does not get much fog at this season. The winter climate is said to be delightful here.

The hills east of Elsinore are the same as noted before and are probably Lower Sonoran.

The sulphur springs & hot baths at Elsinore are the great attraction & in winter the climate is said to be fine & the duck hunting good.

There are lots of ducks on the lake now, mainly cinnamon teal, some godwits & a few mallards. One nest of a teal was found with 2 young & 4 eggs but no young ducks were seen on the lake. A few coots & dabchicks were seen and a flock of about 30 glassy shrikes, some blue herons & a night heron lots of killdeer & several black-necked stilts. These were evidently attracted to the lake in part by the swarms of lake flies breeding in the muddy margins.

Pocket gophers are numerous in the sandy shores of the lake but the few specimens caught do not show any marked local variation.

## Elsinore to Hemet.

July 29 Left Elsinore at 7 A.M.  
and followed up a dry wash, most of  
the way to Perris. The water has been  
taken out of the creek but stands  
in pools here & there. The hills are  
low & rough & in many places a mass  
of granite boulders, but strips of  
flat land make farms here & there.  
The vegetation is mainly 'sages' -  
*Artemisia californica*, *Rhus* *polytricha*,  
& *Stachys* & *Eriogonum fasciculatum*.  
Two juniper trees were seen but  
no amount of upper Sonoran vegetation.  
At Perris Pepper trees, fan palms, mesquites,  
and big Agaves are common, also Turkey  
mullen & tarweed, so I am inclined  
to call it lower Sonoran.

A little east of Perris is an *Atriplex*  
flat & desert like valley. Then  
near Hemet fields of *Abronia*  
make acres of pink carpet over  
sandy fields where the grain has  
been cut. The land is generally  
sandy and rather barren except  
when irrigated.



To Idylwild-

July 30, Took stage to Idylwild at 1:30  
& reached there at 5:30 - 20 miles.  
Hinet is 1600 and Idylwild 5280 feet.

Lower Sonoran reaches to the foothills  
at 2000 feet on north slopes and up the slope  
opposite to about 2500

Upper Sonoran chaparral runs from  
these levels up to about 4500 feet on cold  
slopes and 5500 feet on hot slopes.

Transition zone begins in cold gulches  
at about 4500 feet with *Pseudotsuga* <sup>densa</sup> *macrocarpa*  
and Coulter pine and at on more open slopes  
at 5000 feet with *Pinus ponderosa* & *jeffreyi*  
& Coulter, or at about 5500 feet on south slopes.  
It extends up to about 7700 on north and  
8500 on south slopes.

Located at Idylwild in a tent  
under *Pinus ponderosa* & *jeffreyi* &  
*Quercus californicus* at 5300 feet

July 31. Took most of day for  
making out expense account.

Idylwild

Aug. 172 - Trapped up creek into canyon but caught little.

Aug. 3. Got a horse & started up San Jacinto peak. Climbed the main ridge to 8200 feet, then followed trail north east to creek valley at 7500 feet, then up it to 8500 feet on east side of Marion Mtn. and camped for the night. Set traps from 8500 to 9000 feet in Canadian zone meadows. Slept with only a saddle blanket & had to keep a little fire to keep warm.

Aug. 4. Caught only Microtus & Peromyscus & a Sorex, but left part of traps as I went on over the ridge into Round Valley (Tamoraack Valley) to the Tent kept there by the hotel people during summer. This is at 9000 feet by beautiful Canadian zone meadows. Left my horse & climbed 1800 feet up to San Jacinto peak & back in P.M. A good trail goes to the top & a horse may be ridden to 100 feet of the top.



Found snow banks at 10000 feet and above in many places & top of peak covered 5 snowbanks on cold slope of Jean Peak and rode over one bank of old snow lower down at 9500 feet. There are no meadows or wet places above 9300 feet and the higher slopes are very dry, bare granite gravel which prevents the possibility of many Hudsonian plants. *Pinus murrayana* & *phylicus* reach the peak but for the upper 500 feet are much dwarfed and often prostrate. Quarried & twisted old trunks lie on the ground while the branches reach off from the wind.

The view from the peak is superb, reaching from the ocean to the desert beyond the Salton Sea, from the San Bernardino range to peaks near the Mexican boundary.

Returned to tent, set traps & got my supper & went to bed with plenty of blankets.

Santa Jacinto Mts.

Aug. 5. Caught 2 *Thomomys*, a *Microtus* & *Sorex*. Rode down the creek, North Fork of Tahquitz, about 2 miles to 8500 feet, or to edge of transition zone on the warm slope, then back by the same trail and over the ridge and down to Camp. Set traps in a meadow at 8000 feet for *Thomomys* & *Sorex*.

Aug. 6. Took all day for making up specimens, writing notes and cleaning up from my trip.

Aug. 7. Returned to traps on ridge at 8000 feet and went to top of Tahquitz peak at 8826. Got a *Thomomys* & *Sorex* & some *Peromyscus* colored in were of the zone wasp.

Aug. 8. Shot a *Eutamias merriami* at 5500, this extreme upper limit, & set a few traps. Made up specimens & wrote up notes.



Aug. 9 & 10. - Caught a Weevor  
& wrote on reports.

Aug. 11 - Rode over to Hunt Reservoir  
& then south up the valley to near  
southern end, mapping zones  
on both sides as I went.

Found a big sagebrush valley  
about 10 miles long with lots of desert  
species, *Amphispiza nevadensis*,  
*Chondestes*, *Cyanospiza* etc.

Also jack rabbit tracks & badger holes.

Rode about 30 miles on a hard  
gaited horse & am tired & sore.

At about 4:15 P.M. an earthquake  
shook the camp and ranches, but  
as I was panding along on horseback  
did not feel it. A ranchman  
said it shook his house violently  
& he had telephoned to San Francisco  
& learned that it had broken the  
windows in the bank there.

Aug. 12, Wrote reports & packed specimens

Aug. 13, Staged down to Hemet and got upper limits of the species notes in coming up the mountains.

Corrected zone map & reports accordingly.

Aug. 14, Got a team and drove to Lakeview, and north around the end of San Jacinto Lake and back along the east side of the lake and marshes.

Found many species of water birds and waders and got good zone lists of plants. The desert species are conspicuous north east of the lake where they come down from San Jacinto pass. A few mesquites of both species were seen, also *Chilopsis*, *Atriplex*, *Baccharis*,

Cactus wrens were common & nests seen.

*Dipodomys*, *Perodipus* & *Perognathus* tracks were numerous

Returned to Hemet & set a few traps before dark.

## To Riverside

Aug. 15. Took one P.M. train to Riverside and stopped at Glenwood Hotel. Visited Dr. Attwood in evening & learned much of the country & fauna.

Aug. 16. Mrs. Attwood took us out to the Wilder ranch 7 miles west of town but the wilders had gone to the mts. Got good notes & mapped some country. Took 1:30 train for Redlands & arrived at 4 P.M. Had to wait at San Bernardino for train.

Aug. 17 Took stage for Bluff Lake camp in San Bernardino Mts. & followed up Santa Ana River valley about 10 miles. Then packed our food on horses & mules for about 5 miles up the canyon to fork of Bear Creek then took another old stage <sup>(at 3500 feet)</sup> & came on to Clark's ranch at 4500 feet, then over ridge at 7700 feet, and down to camp at 7500 feet.

The camp is 3 or 4 miles south of Bear Lake. Made lists of plants & birds & mapped zones as I went.



## Bluff Lake

Aug. 18. Sunday, Remained at camp writing up notes, coloring maps most of day & resting up from a very hard camp climb yesterday. Had to walk most of the last half of trip - from 2700 to 7700 feet.

Aug. 19. Trapped around camp.

Aug. 20. Started for San Geronimo Peak and followed the Seven Oaks Trail to where it crosses the Santa Ana River at 5400 then up through Barton Flats to the head of the South Fork of Santa Ana and up a side canyon to Dry Lake at 9000 feet at north base of Grayback (San Geronimo Peak). Did not reach our camp place till dark as we had some 20 miles to walk with considerable climbing. Young Roger S. Palmer went with me and took a horse to carry our blankets & grub & traps but there were no saddle horses available.

## up San Geronimo Peak

Aug. 21. Left camp at 7 am & went up the north east slope of the mountain to the peak at 11480 feet. Found a broad zone of Hudsonian with prostrate timber of murray & flexilis pines and considerable area of Arctic Alpine on N.E. slope of peak with big permanent snow banks. A little scrubby growth goes to the summit on the south side. The Peak is soft light gray granite as are all the range. Great cirques cut into the N.E. slope in two places & glacial valleys extend below with terminal & lateral moraines but not of very recent date. No meadows or moist places were seen above or near timberline and the plants were mostly of dry soil species. A few acres of Bryanthus grow over the north slope from 10000-11000 feet but it is scattered & does not make a complete carpet. There is no place or feed for Pheasants or Ochotona high up & neither occur. Pink snow is abundant. The absence of Piceas & pine squirrels is conspicuous.

Aug. 22 Caught 6 Microtus m. nevadensis  
+ 3 californicus + some other things -  
Broke camp + walked back to  
Bluff Lake, arriving at 4 P.M.  
In evening gave a talk on Geographic  
Distribution to about 100 campers at  
the bonfire.

Aug. 23 - Made up specimens nearly  
all day, wrote up notes and filled  
in gene map.



